# Dyclo-P 50mg

(Diclofenac Potassium) Tablets USP

**Dyclo-P**, the brand of Diclofenac Potassium, is an anti-inflammatory, analgesic and antipyretic agent.

**COMPOSITION:**
Each film coated tablet contains:
Diclofenac Potassium USP....50mg.
Product Specificatin: USP

**ACTION:**
1. **Dyclo-P** is a nonsteroidal anti-inflammatory agent that possesses analgesic and antipyretic activities. It's mode of action, like that of other nonsteroidal anti-inflammatory agents, is not known; however, its therapeutic action is not due to pituitary-adrenal stimulation, but may be from the inhibitory action in the synthesis of prostaglandin by inhibiting enzyme Cyclo-oxygenase.

2. **Dyclo-P** exhibits very marked antirheumatic, anti-inflammatory, analgesic and antipyretic properties and is thus suitable for use in inflammatory and degenerative rheumatic diseases, as well as for the treatment of post-traumatic inflammation and swelling.

**INDICATIONS:**
Inflammatory and degenerative forms of rheumatism, juvenile rheumatoid arthritis, ankylosing spondylitis, osteoarthritis and survical spondyl arthritis.
- Painful syndromes of the vertebral column.
- Non-Article rheumatism.
- Acute attack of gout.
- Painful post traumatic and post operative inflammation and swelling of orthopaedic surgery.
- Painful and inflammatory condition in gynaecology, e.g. Primary dysmenorrhoea and adnexitis.
- As an adjuvant therapy in severe painful inflammatory infection of ear, nose and throat.
- Painful post-operative, inflammatory condition of dental surgery.

**DOSAGE AND ADMINISTRATION:**

**Adults**
The recommended daily dose is 100mg in two divided doses or 150mg in three divided doses. In migraine an initial dose of 50mg should be taken at the first signs

of an impending attack. In cases where relief 2 hours after the first dose is not sufficient, a further dose of 50mg may be taken. If needed, further doses of 50mg may be taken at intervals of 4-6 hours, not exceeding a total dose of 200mg per day. Tablet should be taken with meal or after meal with water.
Tablet should not be taken on empty stomach.

**Children:**
The use of **Dyclo-P** is not recommended in children below 14 years of age.

**SIDE EFFECTS:**
At the start of treatment, patient may complain of epigastric pain, eructation, nausea and diarrhoea, headache, or dizziness. These side effects, however, are usually of a mild nature and generally disappear after a few days despite continuation of the medication. Skin rashes may also occur very occasionally.

**PRECAUTIONS:**
1. In accordance with currently accepted views on the employment of drugs during pregnancy, **Dyclo-P** should not be prescribed in the first three months of pregnancy unless its use is strictly indicated.

2. Patients with gastro-intestinal disturbances or with history of peptic ulcer, as well as those suffering from severe hepatic or renal damage, should be kept under close observation.

**PHARMACEUTICAL PRECAUTIONS:**
1. Protect from heat, light and moisture.
2. Keep all medicines out of the reach of children.
3. Store below 30°C.

**PRESENTATION:**
**Dyclo-P** tablet is supplied in 2x10's blister pack.

*Manufactured by :*
**Indus Pharma (Pvt.) Ltd.**
65/27, Korangi Industrial Area,
Karachi-74900, Pakistan.

02011400013R1