DYCLO

(Diclofenac Sodium)

Injection: 75mg/3ml
Tablets : 50mg, 100mg

Dyclo is chemically Diclofenac Sodium, it is a potent nonsteroidal anti-inflammatory drug (NSAID) with analgesic & antipyretic properties.

COMPOSITION:

Dyclo 50mg Tablet : Each tablet contains 50mg Diclofenac Sodium BP.

Dyclo SR 100 Tablet : Each tablet contains 100mg Diclofenac Sodium BP.

Dyclo Injection : Each 3ml ampoule contains 75mg Diclofenac Sodium BP.

PHARMACOLOGY:

Diclofenac is a very potent inhibitor of prostaglandin synthesis there by relieving pain & inflammation. The drug penetrates rapidly into synovial fluid & peak levels in synovial fluid are achieved 2 hours after peak plasma concentrations.

DYCLO 50mg & DYCLO SR 100 TABLETS:

Dyclo 50mg tablets are enteric coated and are acid resistant while the Dyclo SR 100 tablet is Prolonged released tablet.
The Dyclo SR Tablet has a special matrix that dissolves gradually. This provides a steady, continuous dose of the medication into the bloodstream over 24 hours, allowing for once-daily dosing.

DYCLO INJECTION:

The onset of action Dyclo (Diclofenac) injection is within 16 minutes and the duration of effect is upto 15 hours. The injection should be administered deep into the gluteal muscle in severe arthritic conditions, trauma, orthopaedic manipulations and postoperatively to control pain & inflammation.

INDICATIONS:

Osteoarthritis, Rheumatoid arthritis, Ankylosing spondylitis, Cervical spondylitis, Gout, Capsulitis of shoulder, Frozen shoulder, Lower back pain, Disc lesions, Dental pain, Musculoskeletal pain/strain, Traumatic pain, Synovitis, Tenosynovitis, Bursitis, Dysmenorrhoea, Renal colic and Postoperative pain.

CONTRAINDICATIONS:

Sensitivity to Diclofenac Sodium, Peptic ulceration, Asthma and to patients hypersensitive to aspirin & other prostaglandin synthesis inhibitors.

SIDE EFFECTS:

Rarely mild gastrointestinal upset has been reported. Other rarely observed side effects include headache, fluid retention & peripheral oedema.

PRECAUTIONS:
Severe hepatic, cardic or renal insufficiency & history of peptic ulcer, use in pregnancy should be avoided due to risk of premature closure of ductus arteriosus with prostaglandin inhibitors.

DOSAGE:

Dyclo 50mg tab. : Usual dose is one tablet 2 or 3 times daily or as directed by the physician.

Dyclo SR 100 tab. : Usual dose is one tablet daily or as directed by the physician.

Dyclo Injection 75mg/3ml ampoule:
One ampoule twice a day (after every 12 hours) in acute arthritic conditions, trauma, orthopaedic conditions and to control severe pain after surgery or as directed by the physician. The injection should be administered deep into the gluteal muscle.

Dosage for children:
1 to 3mg per kg daily in divided doses or as advised by the physician. Not recommended for children under 8 years of age.

AVAILABILITY:

Dyclo 50mg Tab. : 20 Enteric coated tablets in alu-alu packing of 10's.
Dyclo SR 100 Tab. : 30 Prolonged Release Tablets in blisters of 10's.
Dyclo Inj. 75mg/3ml : Pack of 50 x 3ml Ampoules.
Pack of 20 x 3ml Ampoules.
Pack of 5 x 3ml Ampoules.

Keep all medicines out of the reach of children.

Manufactured by :
Indus Pharma (Pvt.) Ltd.
65/27, Korangi Industrial Area,
Karachi-74900, Pakistan.

0201140096R5